


DANGER

KEEP OUT OF REACH OF CHILDREN



BLOSSOM BLESSTM



BIOLOGICAL

A PREPARATION OF THE BACTERIUM PANTOEAE AGGLOMERANS P10C FOR USE AS PROTECTION AGAINST FIRE BLIGHT INFECTION OF PIP FRUIT FLOWERS.



Contains: A minimum of 3×10^{10} CFU/g Pantoea Agglomerans P10c in the form of a wettable powder.



NET CONTENTS

300g

GGBLBL/0525

REGISTERED TO: ADAMA New Zealand Ltd
DISTRIBUTED BY: Grochem®, a brand of ADAMA New Zealand Limited
Level 1, 93 Bolt Road Tahunanui, Nelson 7011
PO Box 1799, Nelson 7040 New Zealand
Telephone 0800 447 624 Email: grochem@grochem.com Website: grochem.com

BLOSSOM BLESS

PLEASE READ LABEL COMPLETELY BEFORE USE

May cause an allergic skin reaction. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause long lasting harmful effects to aquatic life.

THIS PRODUCT MUST NOT BE USED FOR ANY PURPOSE OR ANY MANNER CONTRARY TO THIS LABEL.

HANDLING: Read carefully and follow all instructions. Avoid breathing dust, fume, gas, mist, vapours or spray. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. Avoid release to the environment. Wear protective gloves/ protective clothing including eye/face protection and suitable respirator. In case of inadequate ventilation wear respiratory protection.

STORAGE: Keep Frozen. Store in deep freeze at -18°C or below. Remove only what product is required for immediate use, as product that has been opened or thawed cannot be refrozen. When transporting any distance pack in an insulated container and keep cool. Do not heat or microwave to thaw.

DISPOSAL: This product should be disposed of by using as directed on the label. Destroy unwanted product and containers by burning in an appropriate incinerator.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS8409).

FIRST AID: If medical advice is needed, have product container or label at hand.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice/attention.

IF INHALED: If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms, call a POISON CENTRE or doctor/physician.

24-hour Emergency Telephone Number – for advice contact the National Poisons Centre – 0800 POISON (0800 764 766) – or a doctor, immediately.

For specialist advice in an emergency only, call 0800CHEMCALL (24hrs) or 0800 243 6225 In a transport emergency dial 111.

SPILLAGE: Contain spillage with absorbent material. Dispose of waste in a suitable landfill.

TRANSPORT: Do not carry more than 0.5kg on a passenger service vehicle.

SAFETY DATA SHEET: A Safety Data Sheet (SDS) for Blossom Bless is available from Grochem on request or www.grochem.com

WITHHOLDING PERIOD: 1 day

CONDITIONS OF SALE: Grochem®, a brand of ADAMA New Zealand Limited (Grochem) will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by the Statutes applicable to the sale and supply of these goods. To the extent allowed by such Statutes the liability of Grochem is limited to the replacement of the goods or (at the option of Grochem) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Grochem within thirty days of delivery.

Registered pursuant to the ACVM Act 1997, No. P7164. See www.foodsafety.govt.nz for registration conditions.

Approved Pursuant to the HSNO Act 1996, Number HSR001707. See www.epa.govt.nz for approval controls.

Batch:

DoM:

Expiry:

GCBLBL/0525

DIRECTIONS FOR USE:

Blossom Bless is a wettable powder containing the non-pathogenic bacterium *Pantoea agglomerans*. Use Blossom Bless to reduce the incidence of fire blight (*Erwinia amylovora*) on apple and pear blossom. Blossom Bless acts as a protectant, pre-colonising susceptible flower parts and thus preventing the establishment of fire blight causing bacterium *E. amylovora*.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

APPLES – FIREBLIGHT: Blossom Bless acts as a protectant against fire blight infection and must be applied before infection occurs. Application should be made as soon as flowers are open. Repeat applications to maintain beneficial populations of Blossom Bless.

All susceptible flowers should be sprayed not just the target crop.

USE RATES: The rates provided are for high volume spraying to thoroughly wet the crop to the point of run-off. For concentrate spraying adjust the dilution rates accordingly.

TIMING: Always make the first application at 10% flowering

When two applications are made, the second is best made at 50% flowering

If three applications are made, the second is made at 40% flowering and the third at 80% flowering, or prior to a period conducive to rapid bacterial growth, whichever comes first.

NOTE: Numbers of Blossom Bless bacterium will be greatly reduced after frost and re-application should be considered.

COMPATIBILITY: It is recommended that Blossom Bless is applied alone. Blossom Bless can be mixed with sulphur such as Nimbus™, Kumulus® DF, but not lime sulphur.

DO NOT tank mix Blossom Bless with Polyram® DF. Blossom Bless can be sprayed before or after Polyram DF.

DO NOT tank mix Blossom Bless with Streptomycin. Blossom Bless can be sprayed before or after Streptomycin.

DO NOT apply coppers during the flowering period when using Blossom Bless.

BEES: Good bee activity is vital to transfer the Blossom Bless onto newly opening flowers. Bees should be introduced to the orchard before the first application is made and kept in the orchard as long as possible.

Application: Ensure sprayer is in good condition, thoroughly cleaned and well calibrated.

TANK MIXING:

1. Ensure spray tank is clean and free from any chemical residue.
2. Half fill a 5 litre open top container with cold water (do not use hot water).
3. Add contents of this bag, apply lid securely and shake vigorously for 30 seconds.
4. Leave for 5 minutes and once again shake vigorously.
5. Half full the spray tank with water and then pour in the above suspension.
6. Any lumps should be washed through the sieve into the spray tank.
7. Add the balance of the water to the spray tank and apply immediately. Do not mix more than can be applied within 2-3 hours.

DOSE	SITUATION	APPLICATIONS
Low	Orchards with older varieties on older rootstocks.	Two applications of 15g/100L of water.
Medium	Orchards planted with older varieties on dwarf rootstocks.	Two applications of 22.5g/100L of water OR one application of 15g/100L of water followed by one application of 30g/100L of water
High	Orchards planted with pears or newer varieties (eg Pink Lady, Jazz) planted on new and dwarf rootstocks.	One application of 15g/100L of water followed by two applications of 30g/100L of water