READ SAFETY DIRECTIONS BEFORE OPENING





BUDCYTTM

PLANT GROWTH REGULATOR

FOR THE PROMOTION OF BUD GROWTH ON BLIND APPLE WOOD OF NON-BEARING TREES

grochem

Contains: 1g/L Thidiazuron in the form of a ready to use liquid



NET CONTENTS

5L

GBCYT/0422

REGISTERED TO & DISTRIBUTED BY: AgriNova NZ Ltd, 15 Sunlight Grove, PO Box 50-080, Porirua, New Zealand NZ freephone: 0800 447 624 | AUS freephone: 1800 777 068 Email: grochem@grochem.com | Website: grochem.com

BUDCYT™

Budcyt contains Thidiazuron a plant growth regulator for the promotion of bud growth on blind bud sites on non-bearing apple trees. Non-bearing trees being those that will not bear fruit for one or more seasons following application.

CAUTION: Use impermeable gloves when handling. Triple rinse empty containers and dispose of empty containers safely. Avoid contamination of any water supply with the chemical or empty containers. Store in original container, tightly closed, below 23°C and out of reach of children.

FIRST AID If exposed or concerned - contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

BudcytTM is the trademark of ADAMA NZ Limited.
Registered pursuant to the ACVM Act 1997, No P7428. See www.
foodsafety. govt.nz for registration conditions.

24-hour Emergency Telephone Number – for advice contact the National Poisons Centre – 0800 POISON (0800 764 766) – or a doctor, immediately. For specialist advice in an emergency only, call 0800 CHEMCALL (24hrs) or 0800 243 6225. In a transport emergency dial 111.

SHELF LIFE: When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DIRECTIONS FOR USE

It is an offense for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Withholding Period: Do not use after bud break (BBCH07).

APPLICATION:

- · Shake container gently before use.
- · Apply product undiluted.
- · Add no other chemicals.
- Paint the product on the target area with an even coat, using a clean brush.
- Apply to upper and side surfaces of bare wood sections of target branches.

TIMING:

- Application should be made from four weeks before anticipated bud break until bud break on spur sites.
- Ensure sufficient drying time occurs before rain or heavy dew events.
- Best uptake will occur under warm conditions.

CONDITIONS OF SALE: Grochem®, a brand of ADAMA New Zealand Limited (Grochem) will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by the Statutes applicable to the sale and supply of these goods. To the extent allowed by such Statutes the liability of Grochem is limited to the replacement of the goods or (at the option of Grochem) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Grochem within thirty days of delivery.